

**INFLUENCE OF NUTRITIONAL CONDITIONS ON ROOT SYSTEM DEVELOPMENT,
REPRODUCTIVE KIDNEY LAYING AND HARVEST QUALITY**

Kamol Kuchkinovich Shaimanov

Senior Lecturer

Termiz Institute of Agrotechnology and Innovative Development

Ergash Asadullaevich Rakhmonov

Assistant

Termiz Institute of Agrotechnology and Innovative Development

Ilkhom Zhabborovich Boykoraev

Assistant

Termiz Institute of Agrotechnology and Innovative Development

Kholmamat Khurramovich Choriev

Lecturer, 9th Secondary School

ANNOTATION

The scientific article provides experimental material devoted to the study of the influence of the norms of basic nutrients (NPK) and organic fertilization (manure) on the growth and development of the Kishmish Chornyi grape variety. norms of organo-mineral fertilizers - N₁₂₀ P₉₀ K₃₀ kg / ha and N₁₂₀ P₉₀ K₃₀ + 5 t / ha manure.

Keywords: grapes, growth, development, variety, fertilization, harvest.**INTRODUCTION**

Grapes are one of the most valuable dietary and foodstuffs. Grape berries contain up to 30% of easily digestible sugars - glucose and a small amount of sucrose, as well as a large set of organic acids and mineral salts. Grapes are distinguished by a high content of vitamins of groups A, C, P and B. Wine, bekmes, halva, churchkhela, sorbet, grape honey, syrup, jam, marmalade and other valuable nutritious products are prepared from grapes. Grape processing waste is widely used in production. Alcohol, enanthic ester, oil, vinegar, tartaric acid, fodder yeast, enoc dyes and other products are obtained from them[6].

RESEARCH METHODOLOGY

The grape variety Kishmish Cherny was used as an object of study in the experiment. On the experimental plot, fertilizers were applied to a depth of 25-30 cm. The plots were three-row, of which the average was accounting. Each plot was planted with 240 plants. The experiments were repeated three times. Phosphate fertilizers 75% and potash fertilizers 100% were applied in the second or third decade of November. The remaining 25% phosphorus and 100% nitrogen were added to top dressing during the growing season of plants.

Scheme of the Experiment with Fertilization

1. Control - no fertilizer
2. N₁₂₀ kg/ha
3. P₉₀ kg/ha
4. K₃₀ kg/ha
5. N₁₂₀P₉₀ kg/ha
6. N₁₂₀K₃₀ kg/ha
7. P₉₀K₃₀ kg/ha
8. N₁₂₀P₉₀K₃₀ kg/ha
9. N₁₂₀P₉₀K₃₀+5 t/ha manure kg/ha

RESEARCH RESULTS

The complex of external conditions (aeration, nutrition and moisture supply) have a huge impact on the size of the root system, the nature of its placement along the soil horizons and the formation of an active part capable of maximizing the use of nutrients. The interaction of the root system with the environment, its absorption of nutrients from the soil is an active physiological process associated with the vital activity of the whole organism. A large role in this belongs to agrotechnical methods and, first of all, fertilization, which contributes to the creation of a powerful absorbing root surface, which is extremely important for the mineral nutrition of grapes[1, 2, 3, 4].

The roots of the grape bush develop in a large volume of soil and are much larger than the aerial part in length. Therefore, when developing agrotechnical methods for caring for soil and plants, it is necessary to take into account the nature of the distribution of the root system, which is capable of penetrating deep enough, even into the subsoil, which indicates its ability to grow rapidly and strongly branch (5, 6).

Most researchers note that the bulk of small roots is distributed at a depth where the best combination of moisture, aeration and nutrients (40-60 cm).

The growth of the active part of the roots of grapes largely depends on the type of fertilizer. In experiments to study the effect of fertilizer types on the yield and quality of Kishmish black grapes, as well as the growth dynamics of active roots, 9 options were tested: without fertilizers, nitrogen, phosphorus and potassium fertilizers, taken separately, in paired combinations and against the background of complete mineral fertilization.

The greatest effect in the experiment was obtained with the introduction of nitrogen fertilization. With the introduction of phosphorus for one running. meter of root system formed 46 points of growth. In relation to the control, this amounted to 145%. Less active roots are formed when plants are fertilized with potassium alone - 128%. Of the paired combinations, the vine responds best to nitrogen-phosphorus nutrition. The reaction of active roots to a nitrogen-potassium and phosphorus-potassium combination is very close and significantly inferior to a nitrogen-phosphorus mixture. The average number of developed points of growth per 1 meter of root for these combinations is 147, 163 and 143%, respectively, in relation to the control.

Complete mineral fertilization was found to be the best for the formation of absorbing roots. The length of the roots in this variant turned out to be the greatest and reached 1.5 - 2 m.

One-sided feeding with phosphorus or potassium without nitrogen does not give a strong development of absorbing and overgrowing roots for the reason that when mixed with fertilizers with soil particles, the latter are absorbed by it and pass into compounds inaccessible to the plant. In the presence of ammoniacal forms of nitrogen, which have a physiologically acidic reaction, phosphorus and potassium fertilizers dissolve faster and better and are absorbed by the plant in large quantities. (Table 1).

Table 1. Formation of active roots by grapes, depending on the nutrients introduced into the soil

Experience options	Average length active root, mm.	Number of growth points per square meter of root	Sum of growth points	
			PC.	% to control
Control - without fertilizer	0.7	25	515	100.0
N ₁₂₀ kg / ha	0.8	52	863	167.5
P ₉₀ kg / ha	0.8	46	751	145.8
K ₃₀ kg/ha	0.8	40	661	128.3
N ₁₂₀ P ₉₀ kg / ha	0.9	50	819	159.0
N ₁₂₀ K ₃₀ kg / ha	0.8	48	740	143.6
P ₉₀ K ₃₀ kg / ha	0.8	48	758	147.1
N ₁₂₀ P ₉₀ K ₃₀ kg / ha	1.6	59	98	191.0
N ₁₂₀ P ₉₀ K ₃₀ + 5 t/ha manure	1.9	71	1011	196.3

Studies to study the effect of mineral fertilizers and, against their background, an insignificant amount of organic fertilizers (5 t / ha), which is widely used in the world practice of viticulture, has shown that the annual application of mineral fertilizers at the rate of N₁₂₀ P₉₀ K₃₀ contributes to the stable maintenance of the humus content in irrigated gray soils. soils at the level of 1.1-0.6% (Table 2).

Table 2. Dynamics of changes in the content of humus in the soil when fertilizing for grapes, %.

Experience options	Horizons, cm				
	0-10	10-30	30-50	50-70	70-90
Control - no fertilizer	0.9	0.9	0.8	0.5	0,4
N ₁₂₀ P ₉₀ K ₃₀ - 2 consecutive years	1.1	1.0	0.7	0.6	0,4
Control - no fertilizer	0.9	0.8	0.7	0.5	0,4
N ₁₂₀ P ₉₀ K ₃₀ - 3 years in a row	1.1	1.0	0.9	0.7	0,4
Control - no fertilizer	0.8	0.8	0.7	0.5	0.3
N ₁₂₀ P ₉₀ K ₃₀ - 3 years in a row + 5 t / ha manure	1,2	1,2	1.1	0.9	0.5

In the case of systematic non-application of fertilizers, the conditions for humification of the soil deteriorate, and, consequently, the content of total humus in it. This decrease in the soil

horizon of 10-70 cm is 0.1%. With the deepening of the root-inhabited horizon of grapes from 50 to 90 cm, the humus content to the more superficial layers decreases by 1.8-2.2 times, which, naturally, subsequently affects the development of the plant.

The best conditions for stabilizing the humus content in the root layer of grapes, as our experiments have shown, were observed when the full background of mineral fertilizers was applied annually to the soil with the addition of 5 t / ha of manure. In this case, the humus content in the soil in comparison with other variants of the experiment increased by 0.1-0.2%. Moreover, in the soil horizons of 10-70 cm, there was a stable level of humus content over the years of application - 0.9-1.2%. In our opinion, this is due to the creation of favorable conditions in the soil for the activation of the activity of the soil microflora and the processes of soil nitrification.

Studies have also established a positive effect of phosphorus on the formation of generative organs by plants in the wintering eyes of the vine. Under its influence, plants formed up to 69% of fruiting eyes, while with separate feeding of nitrogen and potassium, only 55 and 54%.

With nitrogen-phosphorus and nitrogen-potassium nutrition, the fertility of the eyes increased to 64%, and with complete mineral nutrition, this figure reached 85%, i.e. increased by 16-30%. In addition, a full-fledged mineral background, in comparison with single-element nutrition, stimulates the establishment of plants by more than 1.3-2.2 times more than two ocellous inflorescences, which, as a result, provided a significant overall increase in yield in these experimental options (Table 3).

Table 3 Influence of various types and combinations of fertilizers on the fruitfulness of grape buds

Experience options	Fruitfulness peepholes to the general the number of kidneys %	Glazkov with two inflorescences fruit buds, %	Eyes with one inflorescence to fruit kidneys, %
N 120 kg / ha	55	22	78
P 90 kg / ha	69	33	67
K 30 kg/ha	54	nineteen	81
N 120 P 90 kg / ha	77	37	63
N 120 K 30 kg / ha	64	twenty	80
P 90 K 30 kg / ha	64	19.5	80.5
N 120 P 90 K 30 kg / ha	85	43	37
Control - no fertilizer	49	twenty	80

CONCLUSIONS

1. The annual application of mineral fertilizers at the rate of N 120 P 90 K 30 with the addition of 5 t / ha of manure contributes to an increase in the content of humus in the root horizon of the soil by 0.1-0.2% and its stabilization over the years of application at the level of 0.9-2, one%.
2. When mineral fertilizers are introduced into the plantation of the Kishmish Cherny grape variety and the calculation of N 120 P 90 K 30 kg / ha and N 120 P 90 K 30 + 5 t / ha manure in the

structure of the root system of grapes during the growing season can form up to 59-71 new growth points, and fertility to the total number of buds per individual bush increases by 36% to the control variant.

LITERATURE

1. Abesadze G. The influence of mineral and organic fertilizers on the yield and commercial quality of table grapes.//Tr. 1972. -S. 229-230.
2. Bushin P.M. Fertilization of vineyards. // In the book: Viticulture of Uzbekistan. -Tashkent. 1976. - S. 79-85.
3. Gaprindashvili G.V. The effectiveness of the combined use of organic and mineral fertilizers. // J. Garden and vegetable garden. 1957, No. 7. - 7 p.
4. Garipov R. Growth and productivity of grapes depending on the nutritional regime and load of bushes with eyes. // Intensification of fruit growing and viticulture. -Tashkent. 1987. -S. 45-47.
5. Grigel I.I. The influence of mineral fertilizers on the harvest, the quality of grapes and wine.// Zh. Horticulture, viticulture and winemaking in Moldova. - Chisinau, 1970. No. 9. -WITH. 24-26.
6. Zhurbitsky Z.I. Physiological and agrochemical bases for the use of fertilizers. -Moscow. 1963. -73 s.