

USE OF AMINO ACIDS IN FRUIT AND PEEL OF GARLIC IN MEDICINE

Mamatkulova Ugilkhon

Andijan Institute of Agriculture and Agrotechnologies,
Lecturer at the Department of Physics and Chemistry

Isakov Hayotillo

Andijan State University, Professor of the Department of Chemistry,
Doctor of Technical Sciences

Askarov Ibrokhim

Andijan State University, Professor of the Department of Chemistry,
Doctor of Chemical Sciences

ABSTRACT

The article states that the synthesis of amino acids in the fruits and peel of garlic was determined by the method of Stephen A., Cohen Daviel, comparing the chemical composition of amino acids during processes in the human body. Provides information on the content of amino acids in fruits and peel of garlic

Keywords: assimilation, toxin, hormone, stress ,arthritis ,allergy ,peptide, centrifugation, extraction, glucose, food.

INTRODUCTION

Nature is rich in medicinal plants that have been used in practice since ancient times. Some were discovered through observation, others by studying the experience of using animals and people.

In the 20th century, scientists studied the chemical properties of garlic and proved that the content of garlic is equivalent to penicillin, which kills microorganisms. This can be seen as the use of garlic prevents the growth of fungi and kills parasitic microbes, has a disinfectant effect and is widely used for poisoning. In terms of importance and quantity in the human body, amino acids are second only to water [1]. All currently known amino acids can be of two main types: non-essential and non-essential. Essential amino acids are substances that cannot be synthesized by the body itself and cannot be replaced by other substances, as they regularly enter the body with food.

As for non-essential amino acids, they can be obtained as a result of the synthesis of other nutrients during internal processes, they are not consumed in their pure form. However, these and other amino acids are equally important for the body.[2]

Non-essential amino acids. Non-essential amino acids are synthesized by the body in the process of metabolism, obtained in sufficient quantities from other organic substances. When the need arises, that is, when the amino acids are depleted, the body automatically switches to the mode of creating the necessary amino acids. Non-essential amino acids include the amino acids arginine, alanine, glutamine, glycine, tyrosine, proline, asparagine, serine, and

cysteine. The results of the study show that glutamine is also the most abundant amino acid in garlic.

Glutamine. Essential amino acids can be obtained from many foods, especially greens. This amino acid is involved in building muscles and maintaining their condition. It serves as a source of nutrition for the brain, as well as a source of energy for the nervous system, normalizing its condition and relieving stress [4,6]. In addition, glutamine is able to remove toxic substances from the liver, prevent unwanted breakdown of muscle tissue, strengthen the immune system, and help with arthritis and chronic fatigue. Essential amino acids. Our body cannot synthesize essential amino acids, so almost the only source of them is the food we consume on a daily basis. These include leucine, isoleucine, lysine, methionine, histidine, valine, threonine, and tryptophan.

EXPERIMENTAL PART

The composition and amount of amino acids in the sample was determined in the form of PTC derivatives of amino acids according to the method of Steven A., Cohen D. Isolation of free amino acids. 1 ml (exact volume) of 20% TCA was added to 1 ml of the test sample to obtain precipitation of proteins and peptides from the aqueous extract in centrifuge beakers. After 10 min, the precipitate was separated by centrifugation at 8000 rpm for 15 min. After separating 0.1 ml of the sedimentary liquid, it is freeze-dried. USS analysis of PTK derivatives of amino acids. The synthesis of free amino acid derivatives of FTC (phenylthiocarbomayl) was carried out according to the method of Steven A., Cohen D..

Identification of PTK amino acids is carried out on a Discovery HS C18 75x4.6 mm column on an Agilent Technologies 1200 chromatograph. Solution A: 0.14 M CH₃COONa + 0.05% TEA, pH 6.4, B: CH₃CN. Flow rate 1.2 ml/min, absorbance 269 nm. Gradient %V/min: 1-6%/0-2.5min; 6-30%/2.51-40min; 30-60%/40.1-45min; 60-60%/45.1-50min; 60-0% / 50.1-55min.


RESULTS AND ITS DISCUSSION

Name of amino acids	Garlic	Garlic peel
	Concentration mg/g	
Aspartic acid	0,137844	0,079846
Glutamic acid	0,290147	0,088224
Serene	1,056336	0,275189
Glycine	0,854428	0,133044
Asparagine	0,849209	0,13037
Glutamine	7,259679	0,123217
Cysteine	3,197188	0,466185
Threonine	0,720584	0,355607
Argenin	1,397919	0,280609
Alanine	0,723471	0,141564
Proline	3,724422	0,457229
Tyrosine	0,920939	0,133046
Valine	0,953886	0,410492
Methionine	0,380856	0,016058
Isoleucine	0,293852	0,064873
Leucine	0,539118	0,045519
Histidine	0,324892	0,059806
tryptophan	1,444345	0,14991
Lysine	1,258778	0,01723
Total	26,71584	3,436959

Data File C:\Users\P... \1\Data\DEF_LLC 2022-01-28 09-25-13\002-P1-A2-Chesnok urug'i AAA.D
 Sample Name: Chesnok urug'i AAA

```


=====
Acq. Operator   : SYSTEM                               Seq. Line :    2
Sample Operator : SYSTEM
Acq. Instrument : HPLC                               Location  : P1-A-02
Injection Date  : 1/28/2022 10:03:12                 Inj       :    1
                                                    Inj Volume: 0.000 µl
Different Inj Volume From Sample Entry! Actual Inj Volume : 20.000 µl
Acq. Method    : C:\Users\Public\Documents\ChemStation\1\Data\DEF_LLC 2022-01-28 09-25-13
\AAA.M
Last changed   : 1/28/2022 09:25:11 by SYSTEM
Analysis Method : C:\Users\Public\Documents\ChemStation\1\Data\DEF_LLC 2022-01-28 09-25-13
\AAA.M (Sequence Method)
Last changed   : 1/28/2022 15:35:24 by SYSTEM
(modified after Loading)
    
```



Sample Name: Chesnok pusti AAA

```

=====
Acq. Operator   : SYSTEM                               Seq. Line :    3
Sample Operator : SYSTEM
Acq. Instrument : HPLC                               Location  : P1-A-03
Injection Date  : 1/28/2022 10:40:15                 Inj       :    1
                                                    Inj Volume: 0.000 µl
Different Inj Volume From Sample Entry! Actual Inj Volume : 20.000 µl
Acq. Method    : C:\Users\Public\Documents\ChemStation\1\Data\DEF_LLC 2022-01-28 09-25-13
\AAA.M
Last changed   : 1/28/2022 09:25:11 by SYSTEM
Analysis Method : C:\Users\Public\Documents\ChemStation\1\Data\DEF_LLC 2022-01-28 09-25-13
\AAA.M (Sequence Method)
Last changed   : 1/28/2022 15:35:24 by SYSTEM
(modified after Loading)
    
```



The total amount of amino acids in the studied samples is in the following order: 1 g of garlic in fruit powder (26.7%), 1 g of peel powder (3.44%).

Arrangement of amino acid content (mg) in garlic powder in descending order: Glu (7.26) > Pro (3.7) > Tsis (3.2) > Trip (1.45) > Arg (1.4) > Liz (1.3) > Ser (1.1) > Val (1) > Tyr (1) > Gly (0.9) > Asp (0.8) > Ala (0.7) > Tre (0.7) > Leu (0.5) > Met (0.4) > Fen (0.4) > His (0.3) Iso (0.3) > Gluk-ta (0.3) > Asp (0.2) and etc. [2,3]

The quantitative arrangement of amino acids (mg) in the skin powder is as follows: Cys (0.5) > Pro (0.5) > Val (0.4) > Tre (0.3) > Arg (0.3) > Ser (0.3) > Trip (0.2) > Ala (0.2) > Tyr (0.1) > Asp (0.1) > Gly (0.1) > Gl (0.1) > Gl to-ta (0.1) > Asp (0.08) > Iso (0.06) > His (0.05) > Lei (0.04) > Liz (0.02) > Met (0.02) > Drying (0.01) and so on [2,3,5]

In the sample obtained from garlic powder, the highest amount of glutamine, proline, cysteine, tryptophan and argenin. While the most abundant amino acids in garlic peel powder are cysteine, proline, valine, threonine, and argenine.

Analysis of the results of the analysis of the fruits and peel of garlic makes it possible to choose the ways of using this medicinal plant in the prevention and treatment of diseases associated with the digestive system and the nervous system, for the purpose of general improvement.

CONCLUSION

Thus, the amount of amino acids eaten in garlic and peel, and it was observed that when these amino acids are dried at high temperatures, the powder retains its healing properties. It has also been studied that amino acids play an important role in the processes occurring in the human body by comparing their chemical composition.

REFERENCES

1. Askarov I.R. "Encyclopedia of Medicine" .- Tashkent: Classic word. 2019. 1142 p. (In Uzbek)
2. Olimov N. Q "Author's abstract for the doctoral degree" Development and standardization of technology of drugs based on garlic. " T.2011 (In Uzbek)
3. Usubboev M.U., Olimov N.Q., Aminov S.H. Normurodov A.S. "Development of a tablet drug from garlic powder" // Journal of Pharmacy. -2004. -№3.-P.50-51 (In Uzbek)
4. Mamatkulova O.O., Isakov H., Jamolova H.M. The importance of garlic in the healing of various diseases in humans. // "Folk medicine" magazine. -2021. -№1 (6) .- P.22-24. (In Uzbek)
5. Mamatkulova SA Abstract for the degree of Doctor of Philosophy (RhD) "Classification of individual plants containing lysozyme, pectin, lipids on the basis of chemical composition." F. 2021 (In Uzbek)
6. [Elektronniy resurs: <https://ogorodkand.ru/lechebnye-rasteniya/> -lekarstvenne-svoystva-chesnoka]