

SELECTION OF BREAD WHEAT LINES SUITABLE FOR RAINFED AREAS WITH LOW RAIN IN THE REPUBLIC OF UZBEKISTAN

Dilmurodov Sherzod Dilmurodovich,

Meyliev Akmal Khushvaktovich,
Senior Researchers,

Berdiyeva Feruza Orifovna
Junior Researcher

Southern Agricultural Research Institute,
180100, Karshi City, Kashkadarya Region, Uzbekistan.

ABSTRACT

This article describes the results of experiments on the selection of high-quality lines of bread wheat, as well as the yield of bread wheat in the conditions of rainfed areas, and their transfer to the next stages. The experiment used 33 lines and 2 local check varieties created by local hybridization. Yield of bread wheat lines, 1000 grain vans, grain nature and grain quality indicators were selected for the next stages of selection work by selecting lines higher than the local check varieties.

Keywords: bread wheat, variety, line, 1000 kernel weight, test weight, protein, gluten, vitreous.

INTRODUCTION

Increasing the yield of grain crops for rainfed areas requires the creation of varieties with higher yields of 3-5 c / ha than existing bread wheat varieties and grain quality indicators that fully meet the requirements of a strong wheat class [5, 8, 11, 19, 23, 29, 32, 38, 43, 47].

For this purpose, the evaluation of hybrid lines, which have remained unchanged as a result of selection work, in different soil climatic conditions, on the basis of selection indicators and the selection of the most productive lines and their introduction into production is an urgent task [6, 9, 13, 16, 22, 26, 34, 36, 41, 48, 53].

The high grain weight of 1000 grains in grain formation is a decisive sign of an abundant and stable yield. Lack of moisture in the soil, high temperatures, damage by fungal diseases lead to a decrease in grain weight of 1000 grains [1, 4, 12, 17, 24, 27, 31, 37, 42, 46, 51].

According to scientists, the protein content of wheat grains can range from 8% to 19%, depending on different natural climatic conditions and grain cultivation [2, 7, 15, 20, 21, 28, 33, 39, 44, 49].

According to the experiments of scientists, due to the increase in air temperature in Karshi in 2011 compared to 2010 by 1.4 °C, the protein content of the grain of "Gozgan" decreased by 1.1 %, gluten increased by 2.3 %, yield increased by 9.8 c / ha. showed a decrease. Consequently, high air temperatures and drought conditions have been observed to increase the amount of bread wheat protein and decrease the yield and increase the amount of gluten [3, 10, 14, 18, 25, 30, 35, 40, 45, 50, 52].

RESEARCH METHODS AND MATERIALS

Experimental placement and experimentation were carried out according to the method of phenological observation, calculation and analysis (All-Union Institute of Botany VIR, 1984) and biometric analysis according to the method of the State Variety Testing Commission of Agricultural Crops (1985, 1989). Mathematical and statistical analysis of the experimental results was carried out on the basis of the method of BA Dospekhov (1985). In the research study, the scheme of field experiments was based on the Alpha lattice design of the GenStat 13 program.


Technological quality indicators of winter wheat grain grown in the experimental field "Methodical recommendations for the evaluation of the quality of grain", "Methods of biochemical research of plants", gluten content GOST 13586-1-68, grain glass GOST 10987-76 93, grain nature was studied in comparison with GOST 3040-55, 1000 grain weight in accordance with GOST 10842-89.

The study was conducted in the field experimental field of the Kashkadarya branch of the Research Institute for Grain and Leguminous Crops, located in the territory of Kamashi district, Kashkadarya region. In the experimental field, in the selection nursery of high-yielding, high-quality lines of bread wheat, 35 entry of 2 replication and Gallakor and Ravon varieties were planted as local check varieties. The crop area of each line is 5 m².

RESULTS


One of the necessary indicators characterizing the technological properties of grain depends on the amount of gluten in wheat grain and its properties in the physicochemical composition of the protein that forms mainly gluten in the baking process.

Yields of bread wheat varieties and lines ranged from 1.07 to 3.80 t / ha, while the average yield was 2.92 t / ha. Yield was 3.16 t / ha for local check Gallakor variety and 2.72 t / ha for Ravon variety, while 14 bread wheat lines yielded higher yields than local check varieties.



Picture 1. Higher grain yield lines of bread wheat.

The weight of 1000 grains is one of the main indicators of high productivity. Of the 35 varieties and lines studied, 1000 grains weighed 32.1 to 47.9 g, while the average 1000 grains weighed 40.9 g. A grain weight of 40 grams or more per 1000 grains has a great positive effect on increasing productivity. It was found that the local check grain weight was 38.0 g per 1000 grains, the Ravon variety was 38.9 g, and there were 24 bread wheat lines with higher performance than the local check varieties.



Picture 1. Higher 1000 grain weight lines of bread wheat.

The high grain weight of 1000 can be explained by the fact that in the early stages of selection work the main focus is on grain size. KR18-BW-Sel F5-P-2122, KR18-BW-Sel F5-P-2650, KR18-BW-Sel F5-P-23 lines with a grain weight of 47 grams and more per 1000 grains can be widely used in selection work and recommended in dryland conditions.

Table 1 Productivity and grain quality indicators of bread wheat lines

№	Нав ном	Grain yield, t/ha	1000 grain weight, g	Test weight, g/l	Protein content, %	Grain moisture, %	Gluten content, %	IDK indicators	Grain vitriosity, %
1	Gallakor (check)	3.16	38.0	817.2	16.4	7.2	28.1	87.1	43.8
2	Ravon (check)	2.72	38.9	783.0	14.5	6.8	24.9	85.3	39.5
3	KR18-BW-Sel F5-P-5	2.04	44.2	787.0	15.3	7.1	24.3	79.1	30.8
4	KR18-BW-Sel F5-P-22	2.59	43.3	775.4	14.0	7.0	21.0	86.4	41.0
5	KR18-BW-Sel F5-P-23	3.61	47.9	774.3	14.2	7.0	29.0	92.2	59.8
6	KR18-BW-Sel F5-P-24	2.79	40.4	802.2	14.2	7.2	27.4	83.7	42.8
7	KR18-BW-Sel F5-P-26	3.35	42.9	803.7	15.0	7.1	28.2	87.8	51.5
8	KR18-BW-Sel F5-P-27	2.23	43.3	736.9	14.3	7.1	24.4	91.4	30.8
9	KR18-BW-Sel F5-P-30	3.20	42.6	793.6	15.4	5.8	29.0	94.5	62.8
10	KR18-BW-Sel F5-P-67	3.17	36.6	784.9	14.0	7.1	22.2	75.7	42.8
11	KR18-BW-Sel F5-P-68	3.06	38.4	791.4	14.4	7.0	18.2	80.4	36.3
12	KR18-BW-Sel F5-P-261	2.53	40.0	806.3	14.7	7.2	26.0	83.3	33.3
13	KR18-BW-Sel F5-P-285	3.19	43.9	791.0	14.6	7.2	23.0	92.1	32.0

14	KR18-BW-Sel F5-P-365	2.99	45.0	816.8	15.3	7.3	28.0	74.7	56.3
15	KR18-BW-Sel F5-P-376	2.47	41.4	785.5	14.1	7.9	20.9	83.4	30.8
16	KR18-BW-Sel F5-P-424	3.40	42.1	791.8	15.1	7.3	28.7	83.2	52.3
17	KR18-BW-Sel F5-P-439	2.66	42.7	790.2	15.8	6.9	27.0	75.4	46.0
18	KR18-BW-Sel F5-P-889	2.53	33.0	802.9	14.2	7.0	26.3	79.7	39.8
19	KR18-BW-Sel F5-P-923	2.40	39.7	798.9	13.5	7.5	22.9	94.4	42.8
20	KR18-BW-Sel F5-P-960	3.52	42.9	794.7	15.8	7.3	28.2	88.3	57.3
21	KR18-BW-Sel F5-P-965	3.79	43.4	803.7	15.7	7.2	28.8	80.7	62.8
22	KR18-BW-Sel F5-P-982	3.57	34.8	766.3	14.8	7.1	21.9	101.3	35.5
23	KR18-BW-Sel F5-P-1205	1.07	42.2	789.5	15.2	7.3	20.1	87.6	38.8
24	KR18-BW-Sel F5-P-1212	2.19	32.2	796.3	13.6	6.8	18.9	82.8	21.8
25	KR18-BW-Sel F5-P-1404	2.78	42.1	803.5	13.9	7.4	18.2	84.6	33.3
26	KR18-BW-Sel F5-P-1577	3.32	38.8	743.0	14.2	7.2	16.2	91.4	40.8
27	KR18-BW-Sel F5-P-1579	2.86	38.0	799.1	14.7	7.3	26.2	92.7	40.8
28	KR18-BW-Sel F5-P-1586	3.80	37.0	817.2	14.7	6.8	28.4	81.3	53.3
29	KR18-BW-Sel F5-P-1589	3.03	36.4	803.9	15.2	7.4	28.5	74.3	45.8
30	KR18-BW-Sel F5-P-2122	3.58	47.3	804.5	14.6	7.2	28.6	74.7	54.8
31	KR18-BW-Sel F5-P-2159	3.26	40.8	807.8	14.3	7.4	25.8	85.3	50.8
32	KR18-BW-Sel F5-P-2650	2.80	47.5	805.3	14.2	7.0	28.1	91.5	38.3
33	KR18-BW-Sel F5-P-2825	2.86	43.1	760.8	13.4	7.2	16.3	102.6	23.3
34	KR18-BW-Sel F5-P-2878	3.24	40.1	793.7	14.4	6.8	26.3	92.6	35.0
35	KR18-BW-Sel F5-P-2880	2.60	39.9	808.3	15.4	9.4	30.2	92.6	27.8

Minimum	10.7	32.15	736.9	13.35	5.8	16.2	74.3	21.75
Mean	29.22	40.9	792.3	14.6	7.2	24.8	86.1	42.1
Maximum	38.00	47.9	817.2	16.4	9.35	30.15	102.6	62.75
LSD (0.05)	0.39	1.01	15.06	0.40		1.08		
LSD (0.05), %	1.34	2.46	1.90	2.73		4.35		
CV %	1.6	1.2	0.9	1.3		2.1		

The gluten content of the grain was 28.1% in the local check Gallakor variety and 24.9% in the Ravon variety. It was found that the grain content of gluten was higher in 11 lines than in both local check varieties. KR18-BW-Sel F5-P-2880 had a grain gluten content of 30.2%, KR18-BW-Sel F5-P-30 and KR18-BW-Sel F5-P-23 had a grain gluten content of 29% was found to have reached. The IDK index was 87.1 in the Gallakor variety and 85.3 in the Ravon variety, and in 12 samples it was found to belong to the II class in the range of 85-79 and in 5 samples the IDK index belonged to the I class in the range of 75-74 (Table).

When analyzing the swelling of the studied variety and lines, it was found that the grain viscosity was higher in 12 samples than in the local check Gallakor and Ravon cultivars. The vitreousness of the local check grain variety was 43.8% in the lines KR18-BW-Sel F5-P-30, 62.8% in the lines KR18-BW-Sel F5-P-965, 59.8 in the line KR18-BW-Sel F5-P-23, 8% and 57.3% in the KR18-BW-Sel F5-P-960 line.

CONCLUSION

According to the results of the experiment, 14 lines with higher yields than local check varieties, 22 lines with 1000 grain weight above local check varieties and 40 grams, 28 lines with grain protein content above 14%, 12 lines with grain gluten content above 28% was found to be

present. Bread wheat lines with higher performance than local check varieties were recommended for testing in the next stages of selection and for use in hybridization work.

REFERENCES

- 1) Fayzullayev A. Z. et al. Selection of high-yielding and high-quality lines of bread wheat //INTERNATIONAL SCIENTIFIC AND TECHNICAL JOURNAL “INNOVATION TECHNICAL AND TECHNOLOGY”. – 2020. – Т. 1. – №. 3. – С. 10-14.
- 2) Dilmurodov S. Some valuable properties in evaluating the productivity of bread wheat lines //INTERNATIONAL SCIENTIFIC AND TECHNICAL JOURNAL “INNOVATION TECHNICAL AND TECHNOLOGY”. – 2020. – Т. 1. – №. 1. – С. 60-62.
- 3) Дилмуродов Ш. Д., Зиядуллаев З. Ф. Selection of early and productive lines in preliminary yield trial of bread wheat //INTERNATIONAL SCIENTIFIC AND TECHNICAL JOURNAL “INNOVATION TECHNICAL AND TECHNOLOGY”. – 2020. – Т. 1. – №. 1. – С. 55-59.
- 4) Дилмуродов Ш. Д., Зиядуллаев З. Ф. Юмшоқ буғдойда ўтказилган оддий ва мураккаб дурагайлаш ишлари натижалари //Life Sciences and Agriculture. – 2020. – №. 2-1. – С. 75-79.
- 5) Дилмуродов Ш. Д., Бойсунов Н. Б. Рақобатли навсинаш кўчатзорида юмшоқ буғдойнинг биометрик кўрсаткичларини ўрганиш //Life Sciences and Agriculture. – 2020. – №. 1.
- 6) DILMURODOVICH D. S. et al. Productivity, quality and technological characteristics of bread wheat (*Triticum aestivum* L.) variety and lines for the southern regions of the Republic of Uzbekistan //Plant cell biotechnology and molecular biology. – 2021. – С. 63-74.
- 7) Dilmurodovich D. S. et al. Analysis of yield and yield components traits in the advanced yield trial of winter bread wheat //International journal of discourse on innovation, integration and education. – 2021. – Т. 2. – №. 1. – С. 64-68.
- 8) Dilmurodovich D. S., Bekmurodovich B. N., Shakirjonovich K. N. Winter bread wheat grain quality depends on different soil-climate conditions //International journal of discourse on innovation, integration and education. – 2020. – Т. 1. – №. 5. – С. 377-380.
- 9) Sh K. N. et al. Selection of early bread wheat lines based on studying the time of development //INTERNATIONAL SCIENTIFIC AND TECHNICAL JOURNAL “INNOVATION TECHNICAL AND TECHNOLOGY”. – 2020. – Т. 1. – №. 2. – С. 69-71.
- 10) Dilmurodov S. D., Tukhtayeva U. A. Selection of high-yielding and grain-quality donors of winter bread wheat for irrigated areas //НАУКА И ОБРАЗОВАНИЕ: СОХРАНЯЯ ПРОШЛОЕ, СОЗДАЁМ БУДУЩЕЕ. – 2020. – С. 92-95.
- 11) Dilmurodov S. D., Toshmetova F. N., Fayzullayeva D. Selection of high-quality donor varieties of bread wheat for hybridization //МОЛОДЫЕ УЧЁНЫЕ РОССИИ. – 2020. – С. 55-58.
- 12) Дилмуродов Ш. Д. Подбор исходного материала для селекции пшеницы озимой мягкой для условий Узбекистана на основе изучения хозяйственно ценных характеристик //Аграрная наука. – 2018. – №. 2. – С. 58-61.

- 13) Дилмуродов Ш. Д. и др. Гибридизация в различном направлении и создание гибридного поколения мягкой пшеницы //Иновационное развитие науки и образования. – 2018. – С. 74-77.
- 14) Juraev D. T. et al. To study the heat resistance features of bread wheat varieties and species for the southern regions of the republic of Uzbekistan //European Journal of Molecular & Clinical Medicine. – 2020. – Т. 7. – №. 2. – С. 2254-2270.
- 15) Odirovich J. F., Anvarovich A. O., Dilmurodovich D. S. VALUABLE PROPERTIES AFFECTING THE HIGH-YIELD ELEMENTS OF DURUM WHEAT //INTERNATIONAL JOURNAL OF DISCOURSE ON INNOVATION, INTEGRATION AND EDUCATION. – 2020. – Т. 1. – №. 2. – С. 37-41.
- 16) Dilmurodov S. D., Toshmetova F. N. SELECTION OF EARLY MATURITY LINES IN AGROECOLOGICAL YIELD TRIAL OF BREAD WHEAT //НАУКА, ОБРАЗОВАНИЕ, ИННОВАЦИИ: АКТУАЛЬНЫЕ ВОПРОСЫ И СОВРЕМЕННЫЕ АСПЕКТЫ. – 2020. – С. 41-44.
- 17) Дилмуродов Ш. Д., Бойсунов Н. Б. Отбор продуктивных линий мягких пшениц из гибридного питомника в условиях южного региона Республики Узбекистан //World Science: Problems and Innovations. – 2018. – С. 58-60.
- 18) Дилмуродов Ш. Д. Ценные свойства, влияющие на высокоурожайные элементы мягкой пшеницы //ADVANCED SCIENCE. – 2020. – С. 38-41.
- 19) Дилмуродов Ш. Д. Юмшоқ буғдойнинг маҳаллий маҳсулдор тизмалари селекцияси //Life Sciences and Agriculture. – 2020. – №. 1.
- 20) Дилмуродов Ш. Д., Жабаров Ф. О. Селекция высокоурожайных линий озимой твёрдой пшеницы с высоким качеством зерна //Молодой ученый. – 2019. – №. 31. – С. 34-38.
- 21) Хазраткулова Ш. У., Дилмуродов Ш. Д. Взаимосвязь погодно-климатических условий с продуктивностью и качеством зерна сортов озимой пшеницы //Фундаментальные основы инновационного развития науки и образования. – 2019. – С. 59-61.
- 22) Мейлиев Т. Х., Дилмуродов Ш. Д. Рост и развитие, урожайность и устойчивость к желтой ржавчине сортов в питомнике отбора продуктивных сортов //Приоритетные направления развития науки и образования. – 2019. – С. 130-133.
- 23) Дилмуродов Ш. Д., Бойсунов Н. Б. Селекция местных гибридных линий мягкой пшеницы на юге Республики Узбекистан //Сборник материалов. – 2018. – С. 113-119.
- 24) Дилмуродов Ш. Д., Каюмов Н. Ш. ОЦЕНКА ПРОДУКТИВНЫХ ПОКАЗАТЕЛЕЙ ЛИНИЙ МЯГКОЙ ПШЕНИЦЫ //Вестник науки и образования. – 2020. – №. 17-1 (95).
- 25) Дилмуродов Ш. Д., Бойсунов Н. Б. Юмшоқ буғдой дурагай тизмаларида ҳосилдорликнинг биометрик кўрсаткичларга боғлиқлиги //Иновацион технологиялар. – 2020. – №. 2 (38).
- 26) Dilmurodovich D. S., Shakirjanovich K. N. ANALYSIS OF YIELD AND GRAIN QUALITY TRIALS IN THE ADVANCED YIELD TRIAL OF WINTER BREAD WHEAT //Euro-Asia Conferences. – 2021. – Т. 1. – №. 1. – С. 550-555.
- 27) Dilmurodovich D. S. et al. STUDY OF MORPHO-BIOLOGICAL PROPERTIES AND RESISTANCE TO YELLOW RUST DISEASE OF NEW LINES OF WINTER BREAD WHEAT //InterConf. – 2021.

- 28) Kayumov N. S., Dilmurodov S. D. Selection of heat and drought tolerant varieties and lines of chickpea for rainfed areas //Высокие технологии, наука и образование: актуальные вопросы, достижения и инновации. – 2020. – С. 129-131.
- 29) Shakirjanovich K. N., Dilmurodovich D. S. Analysis of yield and protein content of drought-resistant chickpea lines for rainfed areas //International journal of discourse on innovation, integration and education. – 2021. – Т. 2. – №. 1. – С. 108-111.
- 30) Жураев Д. Т., Дилмуродов Ш. Д. Юмшок бутдойнинг бошоклаш-пишиш даврида иссикликнинг таъсири //Life Sciences and Agriculture. – 2020. – №. 2-2.
- 31) Juraev D. T. et al. Influence of hot dry winds on productivity elements of wheat crop observed in southern regions of the republic of uzbekistan //International journal of applied and pure science and agriculture. ISSN. – 2017. – С. 2394-5532.
- 32) Жураев Д. Т. и др. Влияние суховеев, наблюдаемых в южных регионах республики узбекистан, на продуктивные элементы мягкой пшеницы //Путь науки. – 2017. – №. 2. – С. 84-92.
- 33) Juraev D. T. et al. Influence of hot and dry winds observed in the southern region of the Republic of Uzbekistan on productive elements of bread wheat //The Way of Science. – 2014. – С. 92.
- 34) Juraev D. T. et al. Heritability of Valuable Economic Traits in the Hybrid Generations of Bread Wheat //Annals of the Romanian Society for Cell Biology. – 2021. – С. 2008-2019.
- 35) Дилмуродов Ш. Д. ОЦЕНКА ПРОИЗВОДИТЕЛЬНОСТИ И КАЧЕСТВА ЗЕРНА НОВЫХ МЕСТНЫХ ЛИНИЙ ОЗИМОЙ МЯГКОЙ ПШЕНИЦЫ //Актуальные проблемы современной науки. – 2021. – №. 3. – С. 108-113.
- 36) Dilmurodovich D. S. et al. Selection of large seed and high yielding lines of bread wheat for drought conditions //ACADEMICIA: An International Multidisciplinary Research Journal. – 2021. – Т. 11. – №. 4. – С. 595-606.
- 37) Khushvaktovich M. A., Dilmurodovich D. S. THE CHOICE OF EARLY MATURING LINES OF SPRING BREAD WHEAT FOR IRRIGATED AREAS //НАУКА, ОБРАЗОВАНИЕ, ОБЩЕСТВО: АКТУАЛЬНЫЕ ВОПРОСЫ. – 2021. – С. 30.
- 38) Дилмуродов Ш. Д., Каюмов Н. Ш., Бойсунов Н. Б. ЗНАЧЕНИЕ БИОМЕТРИЧЕСКИХ И ПРОДУКТИВНЫХ ПОКАЗАТЕЛЕЙ ПРИ СОЗДАНИИ ПШЕНИЦЫ С КОМПЛЕКСОМ ЦЕННЫХ СВОЙСТВ //Life Sciences and Agriculture. – 2020. – №. 2-3.
- 39) Хазраткулова Ш. У., Дилмуродов Ш. Д. Оценка жароустойчивости сортов и линий мягкой пшеницы //Наука и образование сегодня. – 2019. – №. 9 (44).
- 40) JURAEV D. T. et al. THE INFLUENCE OF HOT-DRY WIND ON FARM VALUABLE TRAITS OF WHEAT GENOTYPES IN SOUTHERN REGIONS OF UZBEKISTAN //PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY. – 2021. – С. 34-49.
- 41) Файзуллаева Д., Каюмов Н. Ш., Дилмуродов Ш. Д. Лалмикор майдонлар учун нўхатнинг эртапишар тизмалари селекцияси //Молодой ученый. – 2020. – №. 34. – С. 161-163.
- 42) Amanov O. A., Juraev D. T., Dilmurodov S. D. Dependence of Growth Period, Yield Elements and Grain Quality of Winter Bread Wheat Varieties and Lines on Different Soil

- and Climate Conditions //Annals of the Romanian Society for Cell Biology. – 2021. – Т. 25. – №. 6. – С. 5146-5164.
- 43) Dilmurodov S. D. SELECTION OF EARLY MATURITY BREAD WHEAT LINES FOR THE SOUTHERN REGIONS OF THE REPUBLIC OF UZBEKISTAN //Фундаментальные и прикладные научные исследования: актуальные вопросы, достижения и инновации. – 2021. – С. 116-118.
- 44) Dilmurodov S. D. SELECTION OF BREAD WHEAT LINES ON BIOMETRIC INDICATORS HIGH AND LODGING-RESISTANT //НАУКА И СОВРЕМЕННОЕ ОБРАЗОВАНИЕ: АКТУАЛЬНЫЕ ВОПРОСЫ, ДОСТИЖЕНИЯ И ИННОВАЦИИ. – 2021. – С. 65-67.
- 45) Dilmurodov S. D. SELECTION OF EARLY MATURING LINES FROM BREAD WHEAT HYBRID GENERATIONS //НАУЧНЫЕ ИССЛЕДОВАНИЯ МОЛОДЫХ УЧЁНЫХ. – 2021. – С. 81-83.
- 46) Dilmurodov S. D. SELECTION OF F5 HYBRID LINES OF BREAD WHEAT //АКТУАЛЬНЫЕ ВОПРОСЫ СОВРЕМЕННОЙ НАУКИ И ОБРАЗОВАНИЯ. – 2021. – С. 147-149.
- 47) Dilmurodovich D. S. et al. SELECTION OF COLD-TOLERANT VARIETIES AND LINES OF BREAD WHEAT //Наука и образование сегодня. – 2021. – №. 4 (63). – С. 30-33.
- 48) Dilmurodovich D. S., Usmanovna H. S., Sultonovna M. M. SELECTION OF BREAD WHEAT LINES FOR RESISTANT TO THE SOUTHERN HOT CLIMATE CONDITIONS OF THE REPUBLIC OF UZBEKISTAN //Наука и образование сегодня. – 2021. – №. 2 (61). – С. 37-40.
- 49) Dilmurodovich D. S., Usmanovna K. S., Suyarovich I. F. SELECTION OF EARLY MATURING LINES OF WINTER BREAD WHEAT //Вестник науки и образования. – 2021. – №. 4-2 (107). – С. 7-11.
- 50) Egamberdievna A. M., Xudoynazarovich J. B., Dilmurodovich D. S. The effect of sowing time, sowing rates and fertilization rates to field germination of winter bread wheat varieties //ACADEMICIA: An International Multidisciplinary Research Journal. – 2021. – Т. 11. – №. 4. – С. 225-231.
- 51) Дилмуродов Ш. Д., Каюмов Н. Ш., Бойсунов Н. Б. THE VALUE OF BIOMETRIC AND PRODUCTIVE INDICATORS IN THE CREATION OF WHEAT WITH A COMPLEX OF VALUABLE PROPERTIES //Life Sciences and Agriculture. – 2020. – №. 2-3. – С. 50-54.
- 52) Дилмуродов Ш. Д. и др. Сувсизликка чидамли кузги юмшоқ буғдой нав ва тизмаларининг баъзи қимматли хусусиятларини баҳолаш //Молодой ученый. – 2020. – №. 34. – С. 158-161.
- 53) Дилмуродов Ш. Д., Орипов Д. М. Суғориладиган майдонлар учун юмшоқ буғдойнинг F5 авлод дурагайлари селекцияси //Молодой ученый. – 2020. – №. 33. – С. 163-165.
- 54) Xolbazarovich K. K., Sarvarogli M. J., Nikolaevna P. M. Drought and heat tolerance of durum wheat varieties for rainfed conditions of Uzbekistan //ACADEMICIA: An International Multidisciplinary Research Journal. – 2020. – Т. 10. – №. 5. – С. 599-603.