

ANNUAL MATERNAL MORTALITY DETERMINATION FOR LITHUANIA USING THE DOUBLE EXPONENTIAL TECHNIQUE

Smartson. P. NYONI¹

Thabani NYONI²

¹Independent Researcher & Medical Doctor, Harare, Zimbabwe

²Independent Researcher & Health Economist, Harare, Zimbabwe

ABSTRACT

This study uses annual time series data on MMR for Lithuania from 1985 to 2023 to predict future trends of MMR over the period 2024 to 2035. The study utilizes Holt's double exponential smoothing model. The optimal values of smoothing constants α and β are 0.9 and 0.1 respectively based on minimum MSE. The results of the study indicate that MMR will remain very low over the out of sample period. Therefore, we encourage authorities to tirelessly address local causes of maternal mortality and continue routine surveillance of severe maternal morbidity.

Keywords: Exponential smoothing, Forecasting, Maternal Mortality Ratio

INTRODUCTION

Maternal death is a global public health problem. Around 800 women die every day from preventable causes related to pregnancy or childbirth (WHO, 2023). Hemorrhage, hypertensive disorders, and sepsis are responsible for over 50% of maternal deaths worldwide (Khan et al. 2006). In Sub-Saharan Africa, 65% of maternal deaths from hemorrhage occur following delivery (Campbell et al. 2006; Li et al, 1996). Maternal mortality affects low-middle countries more than the developed countries. According to the WHO, at the global level, the maternal mortality ratio in 2020 was estimated at 233 per 100,000 live births (WHO, 2020). However, during the recent years, a significant decline of maternal mortality ratio at the global level was noted, with a reduction of 34.3% between 2000 and 2020, an average rate of reduction of 2.9% per year (WHO, 2023; WBG, 2023; WHO, 2020). The purpose of this paper is to model and forecast MMR for Lithuania using the double exponential technique. The results of the study are expected to highlight future trends of MMR for Lithuania. This will guide policy making, planning and allocation of resources to the maternal and child health program.

LITERATURE REVIEW

Author(s)	Objective(s)	Methodology	Main finding(s)
Mackin et al. (2026)	to understand how levels of MNH care are organized in different country contexts	Scoping review	Three or four levels of MNH facility care were most commonly reported. Linking maternal and newborn facility care to community connections is an important consideration to ensure equitable access to routine and emergency care.

Tsushima et al. (2026)	To determine the long-term incidence of dysglycemia and dyslipidemia after GDM and evaluated whether postpartum changes in body mass index (BMI) independently predicted these outcomes.	Prospective cohort study	Postpartum BMI increase was independently associated with new-onset dysglycemia. No independent predictor of T2DM progression was identified. Dyslipidemia was independently associated with higher pre-pregnancy BMI and multiparity, whereas postpartum BMI change was not independently associated after multivariable adjustment
Yue et al. (2026)	to perform an umbrella review to summarize the current evidence to provide a reference for clinicians and future research	Systematic review	Positive aPLs significantly increased the risk of miscarriage rate in assisted reproductive technology (ART) and were also associated with total fetal loss, PE, intrauterine growth retardation and placental abruption
Nunn et al. (2025)	To explore whether changes in key characteristics among Victorian Aboriginal and/or Torres Strait Islander women contributed to reduced perinatal death rates	Retrospective population-based cohort study of all births in Victoria from 2000–2019	Changes in key maternal characteristics may have contributed to improved perinatal death rates among babies of Victorian Aboriginal and/or Torres Strait Islander women
Haninger-Vacariu et al. (2024)	To describe pregnancy and delivery outcomes in women with Fabry disease	Retrospective cohort-study of women with Fabry disease in Austria using a specific questionnaire and the Austrian Mother–Child Health Passport	The cohort-study in women with Fabry disease showed an increase of pain burden during pregnancies and clearly points to an increased risk for preeclampsia, prematurity, and neonates small for gestational age. With a substantial number of high-risk pregnancies, neonatal outcomes are somewhat worse in Fabry disease than in the general public
Diguisto et al. (2022)	To compare maternal mortality in eight countries with enhanced surveillance systems	Descriptive multi-country population-based study	Variations in maternal mortality ratios exist between high income European countries with enhanced surveillance systems
Morong et al. (2017)	This study was a retrospective cohort study of cases of in-hospital maternal deaths in the Ochsner Health System (Louisiana, United States) and the Queensland Health	-to determine if American and Australian women have a similar rate of preventable maternal death and if the deaths were attributable to the same factors	Women from Louisiana, United States and Queensland, Australia have similar rates of preventable maternal death. No statistically significant factors explained trends in Australian maternal death; American maternal mortality

	System (Australia) from 1995 to 2013		was significantly associated with point of entry into prenatal care, likely influenced by insurance status
Nahidi & Hajifoghaha (2017)	To provide the current situation and trend of MMR in the Eastern Mediterranean region between 1990 and 2015	Systematic review	Based on the collected data, 36 out of 66 studies met the required criteria and were chosen for analysis. All countries of this region showed a downward trend between 1990 and 2015, and this change varied from 16.30% in Yemen to 76.56% in Lebanon. The maternal mortality reduction was 69.88% in Iran in the same time
Schildberger et al. (2017)	-To investigate the personal meanings and experiences of women with disabilities in regard to pregnancy, childbirth and the puerperium	Conducted semi-structured interviews	Participants reported limited acceptance of their life decisions and experienced an environment of discriminatory attitudes. They experienced a lack of support and lack of confidence in their parenting abilities, which negatively influenced their self-efficacy and self-awareness. Violations of personal borders and a feeling of being watched and controlled were reported
Drasutienė et al. (2016)	To describe trends in maternal health in Lithuania during the period 1995–2014 and find causes of maternal mortality	Data was collected from the official statistics database, Medical Data of Births, provided by the Institute of Hygiene Health Information Centre, Vilnius University Faculty of Medicine, and the Vilnius University Centre of Neonatology in 1995–2014	From 1995 to 2014, in Lithuania there were 67 maternal deaths related to pregnancy and childbirth. Hemorrhage and infections during pregnancy, delivery and the postpartum period are considered the most common causes of maternal death
Bouvier-Colle et al. (2012)	To assess capacity to develop routine monitoring of maternal health in the European Union using indicators of maternal mortality and severe morbidity	Analysis of aggregate data from routine statistical systems compiled by the EUROPERISTAT project and comparison with data from national enquiries	Currently available data on maternal mortality and morbidity are insufficient for monitoring trends over time in Europe and for comparison between countries

METHODOLOGY

This study utilizes an exponential smoothing technique to model and forecast future trends of MMR for Lithuania. In exponential smoothing forecasts are generated from the smoothed original series with the most recent historical values having more influence than those in the

more distant past as more recent values are allocated more weights than those in the distant past. This study uses the Holt's linear method (Double exponential smoothing) because it is an appropriate technique for modeling linear data.

Holt's linear method is specified as follows:

Model equation

$$A_t = \mu_t + \rho_t t + \varepsilon_t$$

Smoothing equation

$$S_t = \alpha A_t + (1 - \alpha) (S_{t-1} + b_{t-1})$$

$$0 < \alpha < 1$$

Trend estimation equation

$$b_t = \beta (S_t - S_{t-1}) + (1 - \beta) b_{t-1}$$

$$0 < \beta < 1$$

Forecasting equation

$$f_{t+h} = S_t + h b_t$$

A_t is the actual value of MMR at time t

ε_t is the time varying **error term**

μ_t is the time varying mean (**level**) term

ρ_t is the time varying **slope term**

t is the trend component of the time series

S_t is the exponentially smoothed value of MMR at time t

α is the exponential smoothing constant for the data

β is the smoothing constant for trend

f_{t+h} is the h step ahead forecast

b_t is the trend estimate (slope of the trend) at time t

b_{t-1} is the trend estimate at time $t-1$

Data Issues

This study is based on annual MMR in Lithuania for the period 1985– 2023. The out-of-sample forecast covers the period 2024 – 2035. All the data employed in this research paper was gathered from the World Bank online database.

Findings of the study

Exponential smoothing Model Summary

Table 1: ES model summary

Variable	A
Included Observations	39
Smoothing constants	
Alpha (α) for data	0.900
Beta (β) for trend	0.100
Forecast performance measures	
Mean Absolute Error (MAE)	1.360874
Sum Square Error (SSE)	120.448940
Mean Square Error (MSE)	3.088434
Mean Percentage Error (MPE)	1.121227
Mean Absolute Percentage Error (MAPE)	7.915131

Residual Analysis for the Applied Model




Figure 1: Residual analysis

In-sample Forecast for A




Figure 2: In-sample forecast for the A series

Actual and Smoothed graph for A series




Figure 3: Actual and smoothed graph for A series

Out-of-Sample Forecast for A: Actual and Forecasted Graph




Figure 4: Out-of-sample forecast for A: actual and forecasted graph

Out-of-Sample Forecast for A: Forecasts only

Table 2: Tabulated out-of-sample forecasts

Year	Forecasted MMR
2024	7.7193
2025	7.3720
2026	7.0248
2027	6.6775
2028	6.3302
2029	5.9830
2030	5.6357
2031	5.2884
2032	4.9412
2033	4.5939
2034	4.2466
2035	3.8994

The main results of the study are shown in table 1. It is clear that the model is stable as confirmed by evaluation criterion as well as the residual plot of the model shown in figure 1. It is projected that annual MMR will remain very low over the out of sample period.

RECOMMENDATIONS & CONCLUSION

This study demonstrates the application of the double exponential smoothing method to forecast MMR for Lithuania and findings indicate that MMR will remain very low in the forecast period. Hence, we encourage authorities to address local causes of maternal deaths and continue systematic reporting of severe maternal morbidities.

REFERENCES

1. Campbell O.M., Graham W.J and Lancet Maternal Survival Series steering group (2006). Strategies for reducing maternal mortality: getting on with what works. The lancet, 368(9543), pp.1284-1299.
2. Khan K.S., Wojdyla D., Say L., Gülmezoglu A.M. and Van Look P.F (2006). WHO analysis of causes of maternal death: a systematic review. The lancet, 367(9516), pp.1066-1074.
3. Li X.F., Fortney J.A., Kotelchuck M. and Glover, L.H (1996). The postpartum period: the key to maternal mortality. International Journal of Gynecology & Obstetrics, 54(1), pp.1-10.
4. WBG (2023). Maternal mortality ratio (modeled estimate, per 100,000 live births) 2023. World Bank Open Data. Available: <https://data.worldbank.org>
5. WHO (2020). Maternal Mortality. Available: <https://www.who.int/news-room/fact-sheets/detail/Maternal-mortality>.
6. WHO (2023). Trends in maternal mortality 2000 to 2020: estimates by Who, UNICEF, Unfpa, World Bank Group and UNDESA/Population Division. Geneva: World Health Organization; 2023. Licence: CCBY-NC SA3.0IGO.