

ECOLOGY AND FAUNA OF THE SPECIES BELONGING TO THE FAMILY COLUBRIDAE OF KASHKADARYA REGION

Yangiboyev E. Ch.

PhD Student of National University of Uzbekistan

elmurodyangiboyev20@gmail.com

Rakhimov M. Sh.

Head of Department Zoology of National University of Uzbekistan

raximov.matnazar@mail.ru

ANNOTATION

This article provides informations about the ecology and fauna of the family Colubridae in the Kashkadarya region. Herpetofauna of Kashkadarya region has not been studied scientifically for several decades. Our data were collected in the spring and summer of 2018-2020 and in April-May 2021.

Keywords: reptiles, Folidosis, Colubridae, Typhlopidae, Elapidae, Lcd, Lcr, L, fauna.

INTRODUCTION

Kashkadarya region is located in the southern Uzbekistan, east and west borders are from 39°14'00 "N-66°52'45"E to 38°58'08"N-64°21'58"E and north and south borders are from 39°22'21"N-66°19'20 "E to 38°01'46"N-66°22'02"E. The area is 28.57 thousand km², of which 6.677 thousand km² is arable land, 0.863641 thousand km² is protected area. The average population density is 113 people / km². The main part of the Kashkadarya region is the southern part of the Kyzylkum where are Sandikli desert and Karshi-Nishan steppes (including the lowlands of Guzar district) which covers an area of about 14.5 thousand km², the about 5 thousand km² are foothills and about 9.5 thousand km² consists of mountainous areas[5]. Based on the above data, we are estimated that an average of 80% of the region's area is naturally preserved with excellent fauna and flora. In particular, the literature mentions the information about that 40 species of reptiles are found in the kashkadarya region[1,3].

The distribution and biological characters of reptiles species in the Kashkadarya region was studied by A.M Nikoliskiy in 1915-16, S.K Dalya in 1936, M.V Kaluzhina in 1951, O.P Bogdanov in 1950-60. Our research is of scientific interest, primarily in the fact that scientific research aimed at studying the Reptilia class in the Kashkadarya region was carried out for the last time by V.P Karpenko, X.S Salikhibayev, D.Y Kashkarov, M.M Ostapenko, A.A Petrova, A.Zakirov, N.A Pirnazarov in 1963-65. Then only places of residence of these species was mentioned on the common expeditions of the Institute of Zoology of the Academy of Sciences of the Republic of Uzbekistan[3].

MATERIALS AND METHODS

Field observation and biostatistical analysis of the collected data were mainly used in the study of the Kashkadarya region. The following methods were used.

Selection of the study area: Our research was conducted at 11 points located in the selected 38°52'15.121"N 64°38'36.361"E (250-300 m. above sea level), 39°25'50.281"N 65°26'39.539"E (250-300 m. above sea level), 38°40'27.2532"N 65°35'55.1832"E (350-450 m. above sea level), 38°33'35.4204"N 66°06'27.198"E (400-600 m. above sea level), 38°26'55.425"N 65°53'16.431"E (500-600 m. above sea level), 39°18'44.604"N 66°36'30.5784"E (600-750 m. above sea level), 38°37'07.307"N 66°33'56.996"E (800-1200 m. above sea level), 38°09'35.958"N 66°34'32.181"E (1000-1200 m. above sea level), 39°03'33.798"N 67°12'59.966"E (1800-2200 m. above sea level), 38°41'22.5168"N 67°11'06.6264"E (2100-2400 m. above sea level), 38°49'56.3412"N 67°14'03.904"E (3000-3400 m. above sea level) coordinates across the Kashkadarya region. The selection of these points took into account the biology and ecology of reptiles, the study of natural water bodies, artificial ecosystems, climate and relief effects in the study area, and then the selection of routes. The routes were 4-6 lines that traversed 8-10 km strip crossing the specified coordinates[2,4].

Determining of the Species

Further, the captured reptiles are carried to the base camp, where they are measured and folidosis is done.

Folidosis is a method in which similar species of reptiles are determined by counting certain scales on their body and head. In this case, you must have a measuring tape, ruler, vernier caliper, felt-tip pen, pen and field diary on hand. The last recorded all measurements: L - body length; Lcd-the tail length; Lcr-the length of the head; Scr-the width of the head; Hcr - head height; and also the results of reptile foliosis: Dors - dorsal transverse scales; Vent - ventral scales; An - anal.[2]

Study of Population Distribution in the Area

During the study we studied areas from 120–150 km² (average 135 km²) in each coordinate points. (a total area of 1444.3 km²) It is known that reptiles are spread ruggedly throughout the area. Traces left by individuals are used to study the distribution of the species in the area. for this we need to finding a specific species group (the area around several nests close to each other) and marking the location of the group as the center. we start to observate from the center that there are traces and other nearby nests, as well as traces around them (pieces of rubbish skin and other signs if traces cannot be found). The first area identified during the follow-up will be the active area of the population. Then peripheral area is determined. This area are place where are found individuals of the species or similar marks to this species's marks. And finally, the geographical and ecological boundaries are studied and the full range of the population is determined. During of observation, individuals of a population and its traits may not be present at all in a particular part of the full range, but they may be able to occupy that area in the past or in the future. This method of studying the distribution of species in the study area was calculated using the polygonal convex method[2,4].


Results

During our research, we identified 37 species of reptiles in Kashkadarya region. 7 species are in need of protection listed in the Red Data Book of the Republic of Uzbekistan, 2 species are on the TMXI red list and 1 species is on the CITES red list. 15 of these species belong to the suborder of Ophidia and 4 species are included in the Red Data Book of the Republic of Uzbekistan. 2 of them belong to this family.

During the research, a statistical analysis of 15 species of suborder Ophidia (Serpentes) in Kashkadarya region was divided into 6 families and the following results were obtained. These family are Boidae, Typhlopidae, Elapidae, Viperidae and Colubridae


According to Our research, In Kashkadarya region, 9 species belong to family Colubridae are found and researched. (160 individuals during three years).

The collected materials were analyzed by regions according to the following diagram.



As can be seen from the diagram, members of this family are distinguished by the high number of species and individuals in 8 of the 11 study areas selected in the Kashkadarya region.

At the same time, the population density of individuals was analyzed based on the bio-ecological status of the representatives of this family. The result is that the frequency of meetings of this family in the mountains, hills and deserts throughout the region is given in the diagram below.



Conclusion

In conclusion, our research in Kashkadarya region identified 15 species of Ophidia suborder and they belong to 5 families, 13 genus and studied their ecology and fauna. Consequently, 9 species that are belonging to 6 genus of family Colubridae was found in Kashkadarya region. We have studied only 5% of the total land area of the region, which is 1444.3 km². I think there are still some unexplored species in this area. It should also be noted that today there is a clear decline in the fauna of reptiles due to land development, increasing industrial waste and other similar reasons.

BIBLIOGRAPHY

1. Red book of Republic of Uzbekistan II volume animals. - Toshkent: Chinor ENK, 2019. - p.114-125.
2. Нумеров А. Д., Климов А. С, Труфанова Е. И.//ПОЛЕВЫЕ ИССЛЕДОВАНИЯ НАЗЕМНЫХ ПОЗВОНОЧНЫХ//Издательско-полиграфический центр Воронежского государственного университета 2010г.—42-72

3. Карпенко В., П. Пресмыкающиеся // Экология позвоночных животных Каршинской степи. - Ташкент: Фан, 1967. - С. 129-152.
4. Hester J. M., S. A. Budischak and M. E. Dorcas. 2008. The Davidson College box turtle mark-recapture program: urban herpetological research made possible by citizen scientists. Volume 3, pp. 549–555. Society for the Study of Amphibians and Reptiles, Salt Lake City, Utah.